

Name Calderon

Colombia?



AMERICAN PAD & PAPER CO./HOLYOKE, MASS. 01040

Citadel. NOTEBOOK

50 SHEETS 4X6 INCHES

NO.1372 END OPEN

NO.1373 SIDE OPEN

NARROW RULED



ultimus minor modo
en Colombia 3003

Neurolepis (aristata)

Common on slopes
along rivers. In cloud
forest. Very humid place.
Large clumps ca. 1 m
or more in diameter.
Plants tall, 5-20 m.
Base of the clump covered
by a mat of roots.
No jelly or green stalk
seen. Many new
shoots coming from

buds of the rhizome
Rhizome sympodial with
a lot of branching.

Rhizome develops new ~~shoot~~
shoot which has a very
little developed culm.

with short internodes,
solid, no buds rem-

And no aerial branches

All the new shoots develop
extraxapical, directly

from the rhizome

They are covered by a series of "cabin floors"

The 2-B sand ones

very short, then 1 or 2

with longer with

Developed blade but not

like the shape below.

the leaf ~~is~~ is thicker

188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200.

little demarcation of
blade from outside

Since it is clearly outlined
by legend, it is sym-
metrical. Then come

The true foliage leaves

with the large blade

The sheath of this leaf

is narrow with no

great development. It

continues into the blade

without demerits

It forms a narrow

heart and then gradually

become wider the to

born the blood line

from the store, and
have left for a rest


Wentworth, Long Beach
of L.A. and V.P.R.

on tip and 6 narrow
in 12 cm. at 10 cm.

in bore with pelote

1 leaf measure was:
4.35, by 15.5 cm
in diameter. Midrib
Midrib of leaf not so
thick as in *N. aperta*
leaves are erect and
curve above $\frac{1}{2}$.

Rhizome very short
growing in part, on
top of a rotten tree
trunk.



316

Reserva Natural la Virgen
Páramo de la Virgen - entre
Antio y Papallacta

Growing in so south
slope in páramo repetat-
on eastern slope

Abundant,
2-20 tall.

